



Short Communication

Microneedling with autologous platelet-rich plasma versus topical insulin for treating postacne scars: A split-face comparison

Alpana Mohta¹, Manoj K. Yadav¹, Bhikam C. Ghiya¹

¹Department of Dermatology, Venereology and Leprology, Sardar Patel Medical College, Bikaner, Rajasthan, India.

***Corresponding author:**

Alpana Mohta,
Department of Dermatology,
Venereology and Leprology,
Sardar Patel Medical College,
Bikaner, Rajasthan, India.

dralpanamohta10@gmail.com

Received: 25 January 2023
Accepted: 06 March 2023
Epub Ahead of Print: 04 August 2023
Published:

DOI
10.4103/JCAS.JCAS_20_23

Quick Response Code:



ABSTRACT

Post-acne atrophic scars (PAS) are a common complication of acne, and their treatment can be limited in people of color due to dyspigmentation and scarring risks. This split-face study compared the efficacy of microneedling with topical insulin (TI) on one side of the face with microneedling with platelet-rich plasma (PRP) on the other side for the treatment of PAS. The study included 50 participants with Fitzpatrick skin types III to VI who had never been treated for PAS. Both sides of the face showed significant improvement, with the left side being treated with TI showing a 52% improvement and the right side being treated with PRP showing a 28% improvement. The treatment was well-tolerated, with no significant adverse effects, and no changes in blood glucose levels were observed. The researchers suggest that combining TI or PRP with microneedling can improve PAS treatment, with TI being a superior option due to its ease of use, low cost, and noninvasive nature.

Keywords: Post acne scars, Platelet-rich plasma, Topical insulin, Micro-needling

Postacne atrophic scars (PAS) are a common complication of acne. Peels, platelet-rich plasma (PRP), subcision, dermabrasion, lasers, punch methods, and fillers can be beneficial to PAS. However, many of these can produce dyspigmentation and scarring, limiting their use, especially in people of color.¹ In this population, microneedling is safer and is frequently paired with PRP to boost efficacy, but PRP therapy is pricey and requires more technical expertise.²

This split-face study compared the efficacy of microneedling with PRP on one side of the face (group A), with microneedling with topical insulin (TI) on the other side for the treatment of PAS. We initiated the study after obtaining due institutional ethical approval. The study included 50 participants with Fitzpatrick skin types III–VI who had never been treated for PAS. The following were criteria for exclusion: abnormal blood glucose levels, active acne or infections (herpes, impetigo, seborrheic dermatitis, verruca, molluscum, etc.), pregnancy or lactation, current use of isotretinoin, a history of keloidal propensity, and bleeding diathesis.

Microneedling was done on both sides of the face with a 1.5-mm derma roller with 192 needles on a roller drum. On the left side, we administered 1–2 mL of TI (Human Actrapid Insulin, 40 IU/L solution; Novo Nordisk India Pvt Ltd, Bangalore, India). The right side received 1–2 mL of autologous PRP prepared using standard technique. After 30 min, the face was cleansed, and an ice pack was administered to the treated areas. The procedure was carried out under sterile conditions and with topical anesthetic. The blood glucose levels of

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, transform, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

©2024 Published by Scientific Scholar on behalf of Journal of Cutaneous and Aesthetic Surgery

cases were repeated 2 h before and 2 h after the procedure. There were no harmful consequences observed. Patients were urged to use skin-protective measures. All patients underwent five microneedling treatments monthly and were followed up for 3 months following the last treatment. The Global Acne Scarring System was used to evaluate the treatment's efficacy.

All 50 cases completed the study. The PAS on the left side of the face improved by 52%, and the right side improved by 28% [Figure 1a-d]. A paired *t* test revealed that both sides of the face improved significantly (left side: $P = 0.01$; right side: $P = 0.04$). An unpaired *t* test revealed that both sides of the face improved similarly, although ice pick and boxcar scars responded better to TI therapy. The results were maintained until the last 3-month follow-up.

The treatment was well-tolerated, with no significant adverse effects. Transient erythema, pain, and edema were observed in only a small number of patients, but there was no dyspigmentation or scarring. The researchers did not observe any changes in blood glucose levels.




Figure 1: (a) and (b) Multiple postacne atrophic scars on both sides of the face. (c) and (d) Significant improvement on both sides of the face following microneedling with topical insulin (left side) and platelet-rich plasma (right side).

Type I collagen predominates over type III collagen in a mature PAS.³ Collagen is deposited in a typical lattice pattern following microneedling, and growth factors from PRP can improve PAS healing. Pawar *et al.*⁴ highlighted that to enhance vascular endothelial growth factor, TI activates the phosphatidylinositol 3-kinase/protein kinase B (Akt) pathways. Following TI, the production and maturation of collagen fibers, primarily type III, occur in a basket weave-like pattern (similar to normal skin) rather than a crisscross pattern (similar to scar tissue). As a result, combining PRP or insulin with microneedling can improve PAS treatment.

The study's limitations include limited sample size, a lack of separate assessment of each treatment modality, and a short follow-up period. Employing better dermal delivery systems like tattoo machines could lead to a better depth of drug deposition, as suggested by Ghanian *et al.*⁵

In conclusion, combining TI or PRP with microneedling may increase PAS in people of color while still being safe. However, because of its ease of use, low cost, and noninvasive nature, TI is a superior option to PRP.

Authors' Contributions

Alpana Mohta: Concepts, Design, Definition of intellectual content, Literature search, Manuscript preparation, Manuscript Editing, and Manuscript review. Manoj K. Yadav: Concepts, Design, Definition of intellectual content, Literature search, Manuscript preparation, Manuscript Editing, and Manuscript review. Bhikam C. Ghiya: Concepts, Design, Definition of intellectual content, Literature search, Manuscript preparation, Manuscript Editing, and Manuscript review.

Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent forms. In the form the patient(s) has/have given his/her/their consent for his/her/their images and other clinical information to be reported in the journal. The patients understand that their names and initials will not be published and due efforts will be made to conceal their identity, but anonymity cannot be guaranteed.

Ethical consideration

The study was conducted after obtaining due approval from the institutional ethical board(s).

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

Data availability statement

The data that support the findings of this series are available from the corresponding author with a reasonable request.

REFERENCES

1. Cohen BE, Elbuluk N. Microneedling in skin of color: A review of uses and efficacy. *J Am Acad Dermatol* 2016;74:348-55.
2. Hesseler MJ, Shyam N. Platelet-rich plasma and its utility in the treatment of acne scars: A systematic review. *J Am Acad Dermatol* 2019;80:1730-45.

3. Fabbrocini G, Annunziata MC, D'Arco V, De Vita V, Lodi G, Mauriello MC, *et al.* Acne scars: pathogenesis, classification and treatment. *Dermatol Res Pract* 2010;2010:893080.
4. Pawar M, Singh M. Microneedling with autologous platelet-rich plasma versus microneedling with topical insulin in the treatment of postacne atrophic scars: A simultaneous split-face comparative study. *J Am Acad Dermatol* 2021;84:810-1.
5. Ghanian S, Wambier CG. Response to "Microneedling with autologous platelet-rich plasma versus microneedling with topical insulin in the treatment of postacne atrophic scars: A simultaneous split-face comparative study." *J Am Acad Dermatol* 2021;85:e395-6.

How to cite this article: Mohta A, Yadav MK, Ghiya BC. Microneedling with autologous platelet-rich plasma versus topical insulin for treating postacne scars: A split-face comparison. *J Cutan Aesthet Surg.* doi: 10.4103/JCAS.JCAS_20_23